

The two-domain hypothesis in Beckwith–Wiedemann syndrome: autonomous imprinting of the telomeric domain of the distal chromosome 7 cluster

Flavia Cerrato^{1,†}, Angela Sparago¹, Ines Di Matteo¹, Xiangang Zou², Wendy Dean³, Hiroyuki Sasaki⁴, Paul Smith³, Rita Genesio⁵, Marianne Bruggemann², Wolf Reik³ and Andrea Riccio^{1,*,†}

¹Dipartimento di Scienze Ambientali, Seconda Università di Napoli, via Vivaldi 43, 81100 Caserta, Italy,

²Laboratory of Developmental Immunology, ³Laboratory of Developmental Genetics and Imprinting,

The Babraham Institute, Cambridge CB2 4AT, UK, ⁴Division of Human Genetics, Department of Integrated Genetics, National Institute of Genetics, Research Organization of Information and Systems and

Department of Genetics, School of Life Science, Graduate University for Advanced Studies, 1111 Yata, Mishima,

Shizuoka 411-8540, Japan and ⁵Dipartimento di Biologia e Patologia Cellulare e Molecolare 'L. Califano',

Università di Napoli 'Federico II', Napoli, Italy

Received 18 October 2013; accepted 11 November 2013. doi:10.1016/j.ajhg.2013.11.001

1. $f(x) = \frac{1}{x^2}$ $f'(x) = -\frac{2}{x^3}$ $f''(x) = \frac{6}{x^4}$ $f'''(x) = -\frac{24}{x^5}$

$f(x) = \frac{1}{x^2}$ $f'(x) = -\frac{2}{x^3}$ $f''(x) = \frac{6}{x^4}$ $f'''(x) = -\frac{24}{x^5}$

$f(x) = \frac{1}{x^2}$ $f'(x) = -\frac{2}{x^3}$ $f''(x) = \frac{6}{x^4}$ $f'''(x) = -\frac{24}{x^5}$